

Subjects

- *Indicates book review
- **Indicates correspondence
- †Indicates editor's corner
- ‡Indicates guest editorial

- AA rats, 27-29
- Abstinence
- defining, 150
 - long-term, predictors of relapse in, 640-46
 - in men with severe alcohol problems, 591-98
 - recommended during pregnancy, 19
- Accidental injury, *see* Injuries
- Acetaldehyde
- accelerated production of, 12
 - alcohol toxicity from, 10, 13-14
 - contribution to fetal alcohol syndrome (FAS), 14
 - converted to acetate, via aldehyde dehydrogenase (ALDH) isoenzymes, 133
 - deficiencies in metabolism of, 29
 - ethanol oxidation to, via alcohol dehydrogenase (ADH) isoenzymes, 133
 - and hepatic retention, 14
 - repair of alkylated nucleoproteins impaired by, 13
- Acetaminophen, hepatic injury to alcoholics, 11
- Acetate, acetaldehyde converted to, via aldehyde dehydrogenase (ALDH) isoenzymes, 133
- ACTH release, 17
- Acute hemorrhagic gastritis, 17
- Addiction Research Foundation, *International Profile: Alcohol & Other Drugs*, 246
- Addiction Severity Index (ASI), 56-62, 154-55, 271, 525, 632-33
- European adaptation of, 228
 - suggested modifications to, 60
- ADH isoenzyme (*see also* Gastric ADH)
- in human gastric mucosa, 14
- ADH3 allele frequency, explaining genetic factors, 16
- Admissions, *see* Hospitalizations; Rolling admissions
- Adolescent Diagnostic Interview (ADI), 428
- Adolescent drinkers (*see also* Underage drinking)
- assessing, with Customary Drinking and Drug Use Record (CDDR), 427-38
 - at bush parties, 544-48
 - heavier drinking among, 311-17
 - importance of supervising, 209, 313
 - influences on
 - maternal drinking during and after pregnancy, 292-304, 533-43
 - parents' norms, 33, 311-17
 - peers' norms, 33, 63-70, 208, 311-17
 - and risky sex, 71-77
 - young women, psychoactive substance use disorder (PSUD), and Executive Cognitive Functioning (ECF), 560-67
- Adult drinking culture, *see* Drinking cultures
- Advertising, television, 305-10
- African culture, traditional, rules and regulations for alcohol consumption, 146-51
- Aftercare sessions
- Alcoholics Anonymous (AA) as, 690-97
 - importance of intrapersonal and interpersonal factors in, 161
 - participation in, 276, 521
 - influence on 1-year outcomes, 513-22
 - predictors of, 152-62
- Age differences (*see also* Adolescent drinkers; Children; Older drinkers; Student drinkers; Underage drinking)
- and alcohol toxicity, 14-16
 - and changes in structure of alcoholic beverage preferences, 110
- Aggressive behavior, 563
- inhibiting, 599-607
- AIDS risk behaviors, 154, 156-57
- ALCINVOL, *see* Global scale of alcohol involvement
- Alcohol (*see also* Alcohol effects; Alcoholism)
- administering, intravenously, 259, 266
 - demonizing, 75
 - metabolism of, 9-25
 - calories from, 26
 - first-pass, 14
 - nutritional value of, 26
 - portrayal on prime-time television, 305-10
 - solubility of, 10
- Alcohol abuse, *see* Alcoholism
- Alcohol and other drug (AOD) prevention, 119-20*
- Alcohol-causal hypothesis, 188
- Alcohol-causal-threshold hypothesis, 187
- Alcohol-causal-transient hypothesis, 181, 189
- Alcohol consumption, *see* Drinking
- Alcohol dehydrogenase (ADH), 10
- activity of, 116
 - isoenzymes, ethanol oxidation to acetaldehyde via, 133
- Alcohol Dependence Scale (ADS), 359, 431, 593
- Alcohol Dependence Syndrome, 706
- Alcohol, Drug Abuse, and Mental Health Administration (ADAMHA), 228-29
- Alcohol effects
- castration-like, 15
 - central nervous system (CNS), 293
 - of cocaine administration, 115
 - depressing eating, 27
 - disinhibition, 74
 - effect on ad lib alcohol consumption, 198-206
 - ignoring negative expectancies, 199
 - measuring, 200
 - nondichotomized factor score, 205
 - with drug use, effect of cocaine administration on alcohol consumption, 115-18
 - impairing glucose oxidation, 15
 - intoxication, measuring, 349-50*
 - neuropsychological, 184-86
 - central nervous system (CNS) impairments, 293, 640
 - and organ damage from alcohol abuse, 19
 - pleasurable, 258-69
 - secondary, on co-workers, 608-18
 - sedation, 258
 - on spatial and temporal contrast sensitivity, 163-73
- Alcohol Effects Questionnaire, 410, 412

- Alcohol expectancies (AEs)
 about cognitive impairment by alcohol, 174-79
 activation of, 75
 of adolescent drinkers, and risky sex, 71-77
 defined, 704
 and drinking refusal self-efficacy (DRSE), 704-11
 manipulating of ad lib drinkers', 198-206
 negative (NAEs), 705
 and Theory of Planned Behavior, 409-19
- Alcohol Expectancy Questionnaire (AEQ), 200, 203, 410, 412, 705
- Alcoholic beverages
 beer
 beers per beer-drinking day, 196
 legalization in Iceland, 107-14
 changes in structure of preferences, 109-10
 and age differences, 110
 and drinkers' social profile, 110
 and education, 111-12
 and gender, 110
 and marital status, 110
 and occupation, 112-13
 and urban *versus* rural dwelling, 112
 distilled spirits
 decreases in consumption of, 107
 drinks of spirits per spirits-drinking day, 196
 Gongo, 149-50
 Konyagi, 149-50
 Kindi, 149
 Mapuya, 149
 media portrayal of, 305-10
 Pombe, everywhere in East Africa, 146-51
 Wanzuki, 149
 wine, women getting more of their ethanol from, 195
- Alcoholic brain disease, 92-93
- Alcoholic cirrhosis, 9-25, 92
 association with COL 1A2 locus, 16
 changes in death rates, related to consumption and AA membership, 245-49
 treating with colchicine, 17
- Alcoholic hepatitis, 9, 89, 92
 treating with propylthiouracil, 17
 viral, 16
 B or C, 9
- Alcoholic liver disease, 9-25 (*see also* Alcoholic cirrhosis; Alcoholic hepatitis)
 fatty liver, 10, 16
 fibrosis, 10, 13, 16
 organ damage, 16-19
 toxicity mechanisms of, 10-16
- Alcoholic pancreatitis, 89
 etiology of, 17
 pain relief for, and acetaminophen hepatotoxicity, 11
- Alcoholics Anonymous (AA), 690-97
 affecting alcoholic cirrhosis death rates, 245-49
 as aftercare treatment, 690-97
 barriers to meeting attendance, 694
 compulsory attendance at, 5[†]
 extent of involvement in, 290
 General Service Office, 246
 history of, 469-76
 membership increasing, 245
 as part of patient self-help, 45
 recruiting subjects from, 328
 treatment model, 271
- Alcoholism (*see also* Prevention; Treatment)
 building an identity around, 55
 and child sexual abuse (CSA), 336-48
 chronic, effectiveness of intensive case management for, 523-32
 control mechanisms for, 29
 defining, 373
 and depression, effectiveness of cognitive-behavioral therapy for smokers, 327-35
 developing, 52
 disease model of, 20, 43, 49, 239**, 372
 link with educational achievement, 318-26
 medical disorders of, hepatic, 9-25
 multigenerational, 106
 psychological disorders of
 major depression, 97-106
 suicide, 144
 psychosocial model of, 43-44
 secondary, 98
 undiagnosed, 9
- Alcoholism Rehabilitation Treatment Centers (ARTCs), 470-71
- Alcohol Outcomes Module (AOM), 691-92
- Alcohol oxidation
 by alcohol dehydrogenase, alcohol toxicity from, 10
 producing acetaldehyde, alcohol toxicity from, 13-14
- Alcohol-related diseases, 9-25 (*see also* individual disease conditions)
 hyperlactacidemia, 10
 hyperuricemia, secondary, 10
- Alcohol Research Center Interview, 488
- Alcohol science, birth of, 629[†]
- Alcohol tolerance, acute, 723-30
- Alcohol toxicity
 from acetaldehyde, 10, 13-14
 mechanisms of, 10-16
 role of gastric ADH, 14-16
- Alcohol Use Disorders Identification Test (AUDIT), 420-25, 698-99, 702
- Alcohol Use Inventory (AUI), 272, 593
 global scale of alcohol involvement (ALCINVOL), 272
- Alcohol use trajectories, 647-58
- Aldehyde dehydrogenase (ALDH) isoenzymes, 116
 acetaldehyde converted to acetate by, 133
 individuals deficient in, 133
 protective effect of, 137-38
- Alimentary cancer associated with heavy alcohol consumption, 13
- Alphatocopherol, depleted in cirrhosis patients, 11
- American Association for the Advancement of Science (AAAS), 629[†]
- American Medical Association (AMA), guidelines from, 89, 472
- American Psychiatric Association, *see* Diagnostic and Statistical Manual
- American Society of Addiction Medicine (ASAM), 6[†]
 placement criteria from, 152
- Amino acid sequences, deducing, 14
- γ -aminobutyric acid-A, implicated in alcoholic neurologic complications, 19
- Amphetamine use, in alcoholics with ASPD, 224-25
- Amsterdam Institute for Addiction Research (AIAR), 228
- Amylase, *see* Serum amylase levels
- Analgesics, over-the-counter, 2E1 activation activity towards, 11
- ANA rats, 27-28
- Anemia, megaloblastic, 17-18
- Anesthetics, 2E1 activation activity towards, 11
- Animal studies
 on mice
 bred for high alcohol drinking, 115-16
 cocaine administration lowering alcohol consumption, 115-18
 inherently diabetic, 29
 on nonhuman primates, 10, 16-17
 feeding alcohol to, 16

- of nutritional influences on alcohol intake, 26-28
- of prenatal alcohol exposure, 293
- on rats, bred for high alcohol drinking, 27-30
- Antabuse®
 - administering to prevent relapse, 19
 - contracts, 358-59, 361-62, 365-67, 592
- Anterior cingulate cortex (ACC), smoking lowering activity in, 267
- Antibiotic use, to suppress gastric conversion of urea to ammonia, 17
- 1988 Anti-Drug Abuse Act, 119*
- Antiglutamatergic effects of alcohol, 266
- Antisocial Behavior Questionnaire, 563
- Antisocial personality disorder (ASPD/APD), *see* Psychological disorders of alcoholism
- Anxiety, measuring, 200
- Apparent concomitant motion (ACM), 724-29
- Applied Evaluation Research (AER) portfolio, 120*
- Armed Services Vocational Aptitude Battery, 182
- Ascites, managing, 16
- Assessment of medical disorders, tailoring to individual risk profiles, 94
- Atherosclerosis, protection against high density lipoproteins (HDL), 18
- ATP, *see* Enhanced ATP
- Attrition analyses, 313, 325, 526
- Austin Maze, 184
- Axis I diagnoses, 154, 234-35, 520, 582
- Axis II diagnoses, 227-29, 235
- Azotemia, associated with cirrhosis, 17

- Backward masking, 174
- Barbiturates, increased concentrations when combined with ethanol, 12-13
- Beck Depression Inventory (BDI), 328-30, 332-33, 633
- Beer consumption
 - and increased HDL, 13
 - legalized in Iceland, 107-14
 - rectal cancer associated with low daily intake, 13
- Behavioral counseling (BC), 330-32
- Behavioral impairment, in psychomotor performance, 179
- Behavioral marital therapy (BMT)
 - and behaviorally oriented aftercare following, 372, 591-98
 - and couples relapse prevention sessions following, 357-70
- Behavioral Risk Factor Surveillance System, 122-23**
- Behavior Indicators of Alcoholism Rating (BIAR), 640-46
- Beliefs About Addiction measure, 155
- Benton Visual Retention Test, 184
- Benzodiazepine
 - administering to calm, chlordiazepoxide (Librium®), 19
 - impairment caused by, 267
- Beverages, *see* Alcoholic beverages
- Beverage Strength Rating Scale, 176
- Bexley Maudsley Automated Psychological Screening Test, 184
- Bias, 94, 690-97
- "Big Book," 371, 374
- Binge drinking, *see* Drinking
- Biochemical markers of heavy drinking, carbohydrate-deficient transferrin (CDT), 16
 - in female drinkers, 555-59
- Biologic load, 645
- Block Design Test, 562
- Blood alcohol concentrations (BACs), 724-28
 - determining, 601
 - higher in older drinkers, 15
 - H₂-blocker effect on, 14
 - peripheral, 14
 - in social drinking, higher for women, 14
- Blood alcohol level (BAL), calculating, 349-50*
- Bone fractures, alcoholic proneness to, 15
- Booklet Category Test, 183
- Bootstrap procedure, 575-77
- Breast cancer associated with heavy alcohol consumption, 13
- Breath measuring devices, 165, 176, 200, 217-19, 259, 421, 725
 - performance of, 425
- Brief Michigan Alcoholism Screening Test (BMAST), 420-24, 698-702
- Brief Motivational Counseling, 508
- Brief Symptom Inventory (BSI), 46, 515
 - Global Severity Index, 183
- Bureau of Medicine and Surgery (BUMED), 271
- Bureau of Naval Personnel (BUPERS), 271
- Bureau of the Census, 98
- Bush parties, adolescent drinking at, 544-48

- CAGE questionnaire, 420-26, 698-702
 - defined, 421-22
 - modified, 441-43
- Calories
 - from alcohol, 26
 - tight control of intake, 29
- 1985 Canadian General Social Survey, 78
- Cancers associated with heavy alcohol consumption
 - alimentary, 13
 - breast, 13, 15
 - development of, and synergy with smoking, 13
 - esophageal, 17
 - gastric, 15
 - oropharyngeal, 17
 - rectal, 13
 - respiratory tract, 13
 - statistics on, 93
 - vitamin A deficiency and, 13
- Candies, carrying to prevent relapses, 28
- Carbohydrate-deficient transferrin (CDT) levels, 16
 - in female drinkers, 555-59
- Carbohydrates
 - depressing consumption of, 26
 - increasing, suppresses voluntary drinking, 27-28, 30
- Cardiomyopathy
 - alcoholic, 18, 93
 - higher rates in older drinkers, 89, 94
- Cardiovascular effects
 - large numbers of alcoholic deaths from, 142
 - moderate alcohol use offering protection against, 18
 - and organ damage from alcohol abuse, 18-19
- Care, continuum of, 5-6¹, 152
- β-carotene interaction with alcohol, 13
- Case management, *see* Intensive case management
- Catecholamines, stimulating adrenal medullary secretion of, 17
- Center for Epidemiologic Studies–Depression Scale (CES-D), 401
- Center for Substance Abuse Prevention, 119*
- Center on Alcoholism, Substance Abuse, and Addictions (CASAA), 287-90
- Centers for Disease Control, 455
 - Behavioral Risk Factor Surveillance System, 122-23**
- Central nervous system (CNS) impairment, 293, 534
- Cerebral blood flow (CBF)
 - determining alcohol effects, 258
 - regional, 259-68
- Child Behavior Checklist (CBCL), 401
- Child physical abuse (CPA), and alcoholism, 336-48
- Children, alcohol consumption among, 37-38
- Children of alcoholics (COAs), 387-88, 533-34

- Child sexual abuse (CSA)
and alcoholism, 336-48
committed under influence of alcohol, 72
- Chlordiazepoxide (Librium®), administering to calm, 19
- Chromatography, head-space gas, with flame ionization detector, 116
- Chronic gastritis, 17
- Chronic obstructive pulmonary disease (COPD), alcohol-related, 251, 255
- Chronic public inebriates (CPIs), intensive case management for, 523-32
- Circulating collagenase inhibitor, metalloproteinase and, 16
- Cirrhosis, *see* Alcoholic cirrhosis
- CLABROC, software for receiver operating characteristic (ROC) analysis, 557
- Clinical course of alcohol-related problems, in alcoholic *versus* nonalcoholic dependent drinkers, 581-90
- Clinical research, gap with practice, 125**
- Cocaine
administration, lowering alcohol consumption in mice, 115-18
in alcoholics with ASPD, 225
- Cognitive-behavioral (C-B) therapy
for smokers, with histories of alcohol dependency and depression, 327-35
for substance abuse, evaluating treatment process models, 371-80, 513-22
- Cognitive Behavioral Coping Skills Therapy (CBT), outcomes of, 124**, 504-12, 631-37
- Cognitive Behavioral Mood Management, 330-33
- Cognitive-causal hypothesis, 189
- Cognitive function (*see also* Executive cognitive functioning [ECF])
impaired by alcohol
expectancies about, 174-79
residual effects, 188
in sober social drinkers, 180-90
- Cognitive techniques, 566
- Coherent belief system, affect on staff working environment, 48-49
- Cohesive work setting, 609
- COL 1A2 locus, 16
association with alcoholic cirrhosis, 16
- Colchicine, for treating alcoholic cirrhosis, 17
- Collaborative Study on the Genetics of Alcoholism (COGA) protocol, 485-87, 491, 582
- Collagenase activity
circulating inhibitor, 16
stimulating, 16
- College Alcohol Study (CAS), 122**
- College students, *see* Student drinkers
- Committee on Opportunities in Drug Abuse Research, 237*
- Communities Mobilizing for Change on Alcohol (CMCA), 208
- Community-based programs, 548
- Community-Oriented Programs Environment Scale (COPES), 45
- Comorbidity
of ASPD and depression, 222-26
of FHP and PDs, 495-502
indicating common underlying genetic cause, 97-106
between alcohol dependence and depression, 105
psychiatric, 157
subtyping alcoholics by, 226
- Composite International Diagnostic Interview (CIDI), 228, 234, 318, 421-22
Michigan version of, 320-21
- Computerized Blood Alcohol Calculator (CBAC), 165
- Computers, importance of using for research, 239**
- Conditioned associative-learning task (CALT), 600
- Condom nonuse, and the influence of alcohol, 72
- Conduct disorders (CD), 39, 563
- Consumption of alcohol, *see* Drinking
- Content analysis, 305-8
- Continuity hypothesis, 180, 184
- Continuum of care, 5-6[†], 152
- Contrast sensitivity functions, 164, 166-67
- Controlled Oral Word Association Test, 184
- Convergence, *see* Gender convergence thesis
- Coping factors in mortalities, 144
- Coping Responses Inventory (CRI), 46, 155
- Core Alcohol and Drug Survey, 63-64, 67-69
Long Form, 64
machine scoring of, 64
- Coronary heart disease (CHD), alcohol-related, 250-56
- Cost-effectiveness issues, in U.S. Navy Alcohol Treatment Program, 270-79
- Couples, preventing relapses, after male behavioral therapy, 357-70
- Couples Behavior Questionnaire (CBQ), 359, 592
- Critical fusion frequency (CFF), reduced following alcohol intake, 163
- Cultural changes, 108
- Culture Fair Intelligence Tests (CFIT), 200-201
- Cultures (*see also* Drinking cultures)
traditional African, rules and regulations for alcohol consumption, 146-51
- Current Population Survey (CPS), 672
- Customary Drinking and Drug Use Record (CDDR), psychometric evaluation of, 427-38
- Cytoarchitectonic Area 8, 601
- Cytochrome P-450, 12
- Cytokines, stimulating fibrogenesis, 16
- Deaths (*see also* Suicide)
alcohol-related
in the Lung Health Study, 250-57
underreported, 140
changes in rates of alcoholic cirrhosis patient recoveries, 245-49
from cocaine intoxication, accompanied by alcohol, 115
common causes of, 9
coping factors in, 144
life context in, 144
unnecessary, higher rates of, 6[†]
- Debut of drinking, *see* Drinking
- Decision making, sharing, to improve facility morale, 43-51
- Delinquent behavior, 563
- Delirium tremens*, 19
administering a benzodiazepine to calm, chlordiazepoxide (Librium®), 19
- Denial, 123**
- Dental pain relief for alcoholics, and acetaminophen hepatotoxicity, 11
- Department of Education, Drug Prevention Program in Higher Education, 64
- Department of Health and Human Services, 504
- Department of Veterans Affairs (VA) Medical Centers, 44, 373
- Depression, *see* Psychological disorders of alcoholism
- Depressive-spectrum disease, *see* Psychological disorders of alcoholism
- Dermatologic complications from alcohol abuse, 18
- Descriptive analysis, 571-73
- Designated driving
among student drinkers, 549-54
barriers to use of, 552-53

- Detoxification, outpatient, 5[†]
- Diabetes
 alcohol and, 29
 mice inherently bred to, 29
 noninsulin-dependent, 15
- Diagnostic and Statistical Manual of Mental Disorders (DSM-III)
 criteria for ASPD, 228
 diagnoses of alcohol abuse or alcohol dependence, 183, 223, 641
- Diagnostic and Statistical Manual of Mental Disorders (DSM-III-R), 57, 287, 497, 561
 criteria for adolescent substance use disorders, 428
 criteria for alcoholism (alcohol abuse and/or alcohol dependence), 141, 184, 192, 223, 259, 338, 344, 389, 582, 641, 707
 criteria for ASPD, 227-36
 criteria for major depression, 328
 criteria for personality disorders (PDs), 227-36, 495-502
 criteria for psychoactive substance withdrawal, abuse and dependence, 430-31, 434
 and the Rutgers Alcohol Problem Index (RAPI), 34
- Diagnostic and Statistical Manual of Mental Disorders (DSM-IV), 53
 criteria for alcohol use disorders, 105, 192, 421-22, 670-71
 criteria for psychoactive substance withdrawal, abuse and dependence, 430-31
 definition of abuse, 192
 definition of lifetime alcohol dependence, 97-106, 192
 definition of major depression, 97-106
 definition of trauma, 60
 definition of withdrawal, 193
 overlapping categories within, 477**
- Diagnostic Interview Schedule (DIS), 52, 233-34, 338, 341, 449
- Diagnostic Interview Schedule for Children (DISC), 429, 431, 434
- Diagnostic Interview Schedule—Revised (DIS-R), 287, 328
- Diary reporting, 198, 447-54
- Diet, *see* Nutrition
- Diethylstilbestrol (DES), daughters of mothers who took, 302
- Dilemma test, 34
- Dilinoleoylphosphatidylcholine, 16
- Discharge Information Form (DIF), 44, 373
- Disease concept of treatment (DCT), origins and evolution of, 469-76
- Disease model of alcoholism, 20, 43, 49, 239**, 372
- Diseases (*see also* individual disease conditions)
 resulting from chronic alcohol abuse, 90
 statistics on, 91-92
 sensitive to heavy drinking
 hepatitis, 89
 pancreatitis, 89
- Disinhibition, *see* Alcohol effects
- Disruptive behavior, 563
- Distilled spirits
 consumed more heavily by men, 110
 decreases in consumption of, 107, 718-19
- Dopamine D₂ gene, role in etiology of alcoholism in women, 15
- Dopaminergic systems, mesolimbic, implicated in alcoholic neurologic complications, 19
- Drinker Inventory of Consequences (DrInC), 632-34
- Drinking (*see also* Abstinence; Age differences; Alcohol effects; Alcoholic beverages; Trends)
 ad lib, and manipulating of expectancies, 198-206
 and availability of water, 29
 beliefs about, 67-68
 binge
 changing term to "heavy episodic," 122-24**
 and the five/four measure, 122-23**
 related to DWI (driving while intoxicated) rates, 123**
 and situation in underage drinkers, 213-14
 cessation of
 downward drift to, 144
 manifestations of, 19
 and maximal vulnerability to acetaminophen hepatotoxicity, 11
 curbing, 17
 debut
 consequences of, 32-42
 predictive value of, 32-42
 timing of, 39
 decreases in overall alcohol consumption, 19-20
 hazardous, in the general hospital, 698-703
 heavy
 among adolescents, 311-17
 among East African village dwellers, 151
 among student drinkers, 67-68
 biochemical markers of, carbohydrate-deficient transferrin (CDT), 16
 defining, 182, 671
 diseases sensitive to
 hepatitis, 89
 pancreatitis, 89
 related to job loss, 675-78
 heavy episodic, 122-24**, 311
 inversely related to table sugar intake, 28
 lifetime prevalence of, 383-85
 mechanisms regulating, 29-30
 moderate, 13-14
 cardiovascular benefits of, 18
 defined, 14, 191-97, 638
 rectal cancer associated with, 13
 outcomes, in men with severe alcohol problems, following abstinence, 591-98
 patterns of
 etiology, multifactorial, 133-39
 normal, 32-33
 problem-prone, 32-33
 reduced in Jewish men, association of *ADH2*2* allele with, 133-39
 refusing, 704-11
 social
 advice concerning, 19
 among women, 14
 defining, 182, 185
 effects differing by age, 15
 underreporting, 109
 varying by period (decade), 712-22
 voluntary, 27
- Drinking confederates, 214 (*see also* Influences)
- Drinking cultures
 junior high school students reflecting adult, 381-86
 socialization factors in, 114
 at workplace, 608-18
- Drinking Expectancy Questionnaire (DEQ), 706
- Drinking Habit Questionnaire, 175
- Drinking refusal self-efficacy (DRSE), 704-11
- Drinking Refusal Self-Efficacy Questionnaire (DRSEQ), 706
- Drinking to Cope (DTC) scale, 489
- Drinks
 defining, 65, 108, 671
 Agriculture Research Services' standard, 720
 effects of different approaches to measurement, 191-97

- standard drink units (SDUs), 288
- U.S. standard drinks (USSD), 181-83, 185
- getting larger, 720-21
- presenting, 176
- Driving after drinking (*see also* Designated driving)
 - complexity of, 172
 - related to binge drinking, 123**
 - related to bush parties, 544-48
- Dropping out, *see* Education
- Drownings, correlation with alcoholism, 144
- Drug and Alcohol Program Advisor (DAPA), 271
- Drug and Alcohol Program Treatment Inventory (DAPTI), 45, 374
- Drug Indulgence Index, 430
- Drug Prevention Program in Higher Education, 64
- Drugs, *see* Psychoactive drugs
- Drug Use Chart (DUC), 563
- Drug Use Screening Inventory (DUSI), 428, 563
- DUI/DWI (driving under the influence/driving while intoxicated), 446
 - rates, related to binge drinking, 123**
 - recidivism among, 621-22**
- Dunedin Multidisciplinary Health and Development Study, 302
- Duodenitis, 17
- Dutch Standard Occupation Classification (Standaard beroepen-classificatie), 282
- Eclectic programs, 120*
- Education
 - and drinking, and changes in structure of alcoholic beverage preferences, 111-12
 - dropping out, and risk of alcoholism, 318-26
 - protective effects of, 253-54
- Effects of alcohol, *see* Alcohol effects
- Effects of Drinking Alcohol Scale, 410
- Eighth Special Report to the U.S. Congress on Alcohol and Health, 310
- Electric shocks, administering, 600
- Electroencephalography (EEGs), 350*, 487
- Electronic interviews (EIs), 447-54
- Electro-oculographic (EOG) techniques, 725
- Employee assistance programs, compulsory, 5†
- Employee Health and Performance in the Workplace questionnaire, 610
- Endocrine pathology, and organ damage from alcohol abuse, 17
- Energy value of ethanol, 10, 26
- Enhanced ATP
 - breakdown of, acetate-mediated, 10
- Environmental predictors of alcoholism risk, 485-94
- Epidemiological Catchment Area studies, 222-23, 318
- Epi-info software, 147
- Esophagitis, 17
- Ethanol
 - energy value of, 10
 - psychoactive properties of, 10
- Ethanol-exacerbated lactase deficiency, 17
- Ethnic differences
 - defining, Hispanic, 287, 423, 660
 - in drinking patterns over the life course, 568-80
 - in gastric ADH, 14
 - Jewish drinkers deficient in ALDH activity, 133
 - in treatment utilization and outcome, Hispanics *versus* non-Hispanic whites, 286-91
 - trend to increased alcohol consumption
 - among blacks, 659-68
 - among Hispanics, 659-68
- Ethno-linguistic differences, in Ontario, Canada, 83
- Euphoria, *see* Alcohol effects
- EuropASI interview, 228, 230
- Event-related potentials (ERPs), 183-84
 - in social drinkers, 184-88
- Executive cognitive functioning (ECF)
 - in female adolescents with psychoactive substance use disorder, 560-67
 - impaired, may facilitate aggression, 560
- Expectancies, *see* Alcohol expectancies
- Expected Type of Effect Scale, 175
- Exploratory analyses, 595-96
- Eysenck Personality Questionnaire (EPQ), 200
- Family Environment Scale (FES), 134-35
- Family history (FH) of alcoholism
 - effects of
 - on comorbidity for alcohol dependence and depression, 97-106
 - gender differences in, 97-98
 - disruptive, delinquent and aggressive behavior in female adolescents with a PSUD, 560-67
 - family history positive (FHP) *versus* family history negative (FHN) dichotomy, 99-100, 486-89
 - saturation, 100-104
- Family Tree Questionnaire for Assessing Family History of Drinking/Drug Abuse Problems (FAMTREE), 537
- Fasting, synergy with acetaminophen hepatotoxicity, 11
- Fatal Accident Reporting System (FARS), 465
- Fatty acid synthesis promoted, 26
 - by NADH, 10
- Federal Communications Commission (FCC), 305
- Federal Trade Commission (FTC), 305
- Feedback, role in formation of acute tolerance to alcohol, 723-30
- Feighner Criteria, 223
- Female drinkers, *see* Women
- Feminization, hallmark of cirrhosis, 17
- Fetal alcohol effects (FAE), 620*
- Fetal alcohol syndrome (FAS), 350-52*, 534, 620*
 - acetaldehyde contribution to, 14
 - rodent models of, 352*
- Fibrosis, 10, 13, 16
 - perivenular, 16
 - preventing with polyenylphosphatidylcholine, 17
- First pass metabolism (FPM), gastric, 14
- First-time admissions, *see* Hospitalizations
- Five/four measure of binge drinking, 122-23**
- "5 or more" drinking, 212-13
- Follow-up Information Form (FIF), 514-15
- Food and Agriculture Organization (FAO), 150
- Four Factor Index of Social Status, 563
- France, drinking patterns in, 19
- Francophone/Anglophone differences in alcohol use, 78-88
- Fraternities, *see* Student drinkers
- Free hydroxy-ethyl radical generation, 11
- Free radical-mediated toxicity, promoted by acetaldehyde, 14
- GABA mimetic effects of alcohol, 266
- Gamma-glutamyl transferase (GGT), 700
 - in social drinkers, 188
- Gamma glutamyl transpeptidase (GGTP), as a biological marker, 555-59
- Gastric ADH
 - ethnic differences in, 14
 - inhibition of, 14
 - role in alcohol toxicity, 14-16

- Gastric first pass metabolism (FPM), 14
- Gastrointestinal complications, and organ damage from alcohol abuse, 10, 17
- Gastrointestinal (GI) bleeding, 17
- Gender convergence thesis, 33, 39
- Gender differences (*see also* Men; Women)
in alcohol-related diabetes, 29
and alcohol toxicity, 14-16
and changes in structure of alcoholic beverage preferences, 110
in drinking patterns over the life course, 568-80
vulnerability thesis, 33
- General Background Questionnaire (GBQ), 496
- Genetic-causal hypothesis, 189
- Genetic factors
COL 1A2 locus, 16
defects in aldehyde dehydrogenase, 19
matrilineal heritability of depressive-spectrum disease, 106
reduced alcohol consumption by Jewish men, association of *ADH2*2* allele, 133-39
- Genetic heterogeneity, comorbidity indicating common underlying cause, 97-106
effect stronger in women, 97-98
- Gestalt Closure Test, 184
- Global scale of alcohol involvement (ALCINVOL), 272, 275
- Global Severity Index, 183, 273
- Gluconeogenesis, inhibition by alcohol, 29
- Glucose homeostasis, alcohol-induced alterations in, 30
- Gongo, 149-50
- Go/no-go tasks, 562
- Gouty attacks, reasons for, 10
- Governmental alcohol policies, *see* Policy issues
- Graduated frequency (GF) measure, 34
- Graduate Licensing Study (GLS), 544-47
- Granulocytopenia, 18
- Greek societies, *see* Student drinkers
- Group cohesion
as buffer, 609
and influence of workplace drinking and stress, 608-18
- GSH, *see* Hepatocellular GSH
- Halstead-Reitan Neuropsychological Battery, 184
- Hamilton Rating Scale for Depression (HRSD), 328-29, 332-33
- Hangover effect, 181
- Hanil Alcohol Insight Scale (HAIS), 52-55
- Harm reduction, 619-20*
- Hazardous behaviors, willingness to engage in when little impairment expected, 179
- Headache pain relief for alcoholics, and acetaminophen hepatotoxicity, 11
- Health and Daily Living Form (HDSL), 515
- Health care work environments, in treatment institutions, 43-51
characteristics of, 43
patients' improvement during treatment, 44
patients' involvement in, 44
- Healthier People Health Risk Appraisal, 272
- Health providers, 5-7†
- Heavy drinking, *see* Drinking
- Heavy episodic drinking, *see* Drinking
- Heckman selection model, 691
- Helicobacter pylori* infection, contributing to hepatic encephalopathy, 17
- Hematologic disorders, and organ damage from alcohol abuse, 17-18
- Hematopoiesis, ethanol toxicity and, 18
- Hepatic coma and precoma, 17
- Hepatic encephalopathy
helicobacter pylori infection contributing to, 17
spontaneous, 17
- Hepatic phosphatidylcholine, 16
- Hepatic retention, and acetaldehyde, 14
- Hepatitis, *see* Alcoholic hepatitis
- Hepatocellular GSH
depletion in steady state levels of, 11
promoted by acetaldehyde, 13-14
precursors of, 12
synthesis inhibited, 11-12
- Hepatoprotective agents, 17
- High density lipoproteins (HDL)
increased by moderate alcohol use, 18
protection against atherosclerosis, 18
- Hispanics, treatment utilization rates of, *versus* non-Hispanic whites, 286-91
- HIV high-risk behaviors, 154
- Hollingshead Four Factor Index of Social Status, 563
- Home Environment Interview, 341
- Homelessness, 524
- Homeostasis, redox, 10
- Homocysteine levels, 18
- Hormones, and alcohol toxicity, 14-16
- Hospitalizations
alcohol-related, 9
compared to myocardial infarction-related, 9
emergency room population, alcohol-related, screening, 420-26
first-time admissions, 89-96
intensive care unit admissions, alcohol-related, 9
- Hospitals, hazardous drinking in, screening for, 698-703
- H₂-blocker effect, on blood alcohol concentrations, 14
- Human gastric mucosa, ADH isoenzyme in, 14
- Human Immunodeficiency Virus (HIV) infection danger and alcohol use among adolescents, 71-77
reducing risk in innovative treatment programs, 121*
- Humanism, inadequate to sustain staff environment, 49
- Hyperlactacidemia, 10
- Hyperparametric analysis
defined, 569-70
of drinking patterns over the life course, 568-80
- Hyperuricemia, secondary, 10
- Hypogonadism, hallmark of cirrhosis, 17
- Hypothalamic-pituitary-adrenal axis, alcohol activation of, 17
- Important People and Activities Scale (IPA), 489
- Impulsivity, 501
- Incest victims, 342
- Increased Riskiness scale, 72
- Individual risk profiles, tailoring assessment of medical disorders to, 94
- Inferential analysis, 571, 573-76
- Influences
on adolescent drinkers, 33, 208
maternal drinking during and after pregnancy, 292-304, 533-43
on student drinkers, 69
- Information
garnering, from computer-aided research, 239**
processing, affected by alcohol, 174
- Informed consent, 147, 448
- Injuries, alcohol-related skiing, and sensation seeking, 216-21
- Inner-city children, *see* Socioeconomic status (SES)
- Inpatient treatment
diagnosis preceding, 94
importance of, 5-6†

- keeping option open for some patients, 5[†]
 move to abolish, 5-7[†], 125**
- Input/output placement, random, 6[†]
- Insight and Treatment Attitude Questionnaire, 52
- Insight-evaluation instruments, Hanil Alcohol Insight Scale (HAIS), 52-55
- Institute of Medicine
 Committee on Opportunities in Drug Abuse Research, 237*, 351*
 definition of aftercare, 690
- Intake Information Form (IIF), 44, 373, 514-15
- Intellectual competence, 182
- Intensive case management, for chronic inebriates, 523-32
- Intensive outpatient program (IOP), 153-55, 157, 159-60
- Interactive voice response (IVR) system, 448
- International Classification of Diseases, Version 9 (ICD-9)
 coded discharge diagnoses (ICD-9-CM), 90-91, 93, 373, 514
 coding for cause of death, 141, 251
- International Classification of Diseases, Version 10 (ICD-10), diagnoses, 122**, 229, 420-25
- International Personality Disorder Examination (IPDE), 228-29, 232
- International Profile: Alcohol & Other Drugs*, 246
- Interrater reliability, 54, 433
- Interstimulus interval (ISI), short, 174
- Interviews, for self-reported alcohol use, 280-85
- Intoxication, *see* Alcohol effects
- Intravenous bolus method, 259
- IQ decrements, 534
- Japan National Institute of Mental Health, 382
- Jewish drinkers
 deficient in ALDH activity, 133
 ultra orthodox, declining study participation, 133
- Jewish people, sensitive to heavy drinking, 137
- Job loss, and alcohol misuse, 669-80
- Journal of Studies on Alcohol*, 122**, 473
 authoring for
 need for careful computer-aided research, 239**
 use of term *binge*, 124**, 621**
- Journal of the American Medical Association*, 123**
- Karolinska Scales of Personality (KSP), 489
- Kaufman Hand Movement Test, 184
- Ketosis, alcohol-induced, 10
- Khavari Alcohol Test (KAT), 706-7
- Kiddie-Schedule for Affective Disorders and Schizophrenia (K-SADS), 428
 Expanded (K-SADS-E), 561, 563
- Kindi, 149
- Konyagi, 149-50
- Korean drinkers, beliefs about drinking among, 52-55
- Korsakoff's disease, 180
- Lactase, ethanol-exacerbated deficiency in, 17
- Latent content analysis, 308-10
- Latent Growth Modeling (LGM), 656
 multilevel, 399-408
- Latent messages, about alcohol, portrayal on prime-time television, 305-10
- LEAD (longitudinal, expert, all data) standard, 235
- Legal issues in alcohol, forensic, 350*
- Lexus/Nexus news database, 122-23**
- Libido losses, hallmark of cirrhosis, 17
- Librium®, administering to calm, 19
- Life context in mortalities, 144
- Life Experiences Survey (LES), 489
- Life Stressors and Social Resources Inventory (LISRES), 515
- Lifestyle Choices Survey (LCS), 534
- Lifetime aggression, 565
- Lipid oxidation, opposed by NADH, 10
- Lipid peroxidation, 12
- LISCOMP software, 35-36
- Liver disease, *see* Alcoholic liver disease
- Liver transplantation, 17
- Logical regression analyses, 312, 541
- Log-likelihood tests, 579
- Longitudinal studies, *see* Studies
- Lundby project, 140-45
- Lung Health Study, 250-57
- MacAndrew Alcoholism Scale (MAC), 200, 202, 205, 439-45, 489-90
- Macronutrient intake patterns, 28
- Magnetic resonance imaging (MRI), 350*
- Mail questionnaires, for self-reported alcohol use, 280-85
- Major depression, *see* Psychological disorders of alcoholism
- Mallory-Weiss lesions, 17
- Malnutrition, cause of alcohol toxicity, 10
- Manifest content analysis, 305-8
- Mapuya, 149
- Marijuana, in alcoholics with ASPD, 224
- Marital Adjustment Test (MAT), 359, 361-62, 592
- Marital and family therapy (MFT), 357
- Marital outcomes, in men with severe alcohol problems, following abstinence, 591-98
- Marital status (*see also* Couples)
 affect on drinking patterns, 570, 579
 changes in, and developmental trajectories of alcohol use in young adults, 647-58
 and changes in structure of alcoholic beverage preferences, 110
 married drinkers, under greater stress, 110
- Marital Status Inventory, 359
- MATCH, *see* Project MATCH
- Maternal drinking
 during and after pregnancy, 292-304, 351-52*, 533-43
 effect on breast-feeding, 351*
- Matrilineal heritability, of depressive-spectrum disease, 106
- Mean corpuscular volume (MCV), 700
 increases in, 17-18
- Media, television, prime-time, portrayal of alcohol on, 305-10
- Mediation analyses, 595
- Medical disorders coexisting with alcoholism, 6[†]
 assessment of, tailoring to individual risk profiles, 94
 hospitalizations for, first-time admissions, 89-96
 osseous abnormalities, 15
 pathophysiology of, 9-25
 severity of, 9, 15
- Megaloblastic anemia, 17-18
- Mellanby effect, 721
- Men
 feminization in, hallmark of cirrhosis, 17
 heavier consumption of distilled spirits, 110
 hypogonadism in, hallmark of cirrhosis, 17
 paternal drinking, and paternal absence, development of problem behaviors in boys, 387-98
 testosterone effects, suppressing hepatic ADH activity, 14-15
- Menstrual cycle disorders, 14-15
 hallmark of cirrhosis, 17
- Messenger RNA (mRNA), 16
- Metabolism, *see* Alcohol
- Metalloproteinase, and circulating collagenase inhibitor, 16

- Metadone**
 innovative programs based on, 120*
 and long-term *versus* short-term ethanol administration, 12
- Mice**
 cocaine administration lowering alcohol consumption, 115-18
 inherently diabetic, 29
- Michigan Alcoholism Screening Test (MAST)**, 135, 200, 283, 340-42, 359, 439-45, 593, 600, 698-702
- Microsomal ethanol oxidizing system (MEOS)**
 alcohol toxicity from, 10-13
 induced by chronic ethanol consumption, 10-12
- Mini-Mental State Examination**, 186
- Minnesota Model treatment**, 372
- Minnesota Multiphasic Personality Inventory (MMPI)**, 216, 439, 471, 641-44
- Minnesota Twin-Family Study (MTFS)**, 222-23
- Mitochondrial inefficiency, secondary to chronic ethanol consumption**, 10
- Models**
 multiple group structural equation, 35, 39
 treatment, evaluating, 371-80
- Moderate drinking**, *see* Drinking
- Modified PTSD Symptom Scale (MPSSR)**, 60
- Monitoring the Future project**, 122**
- Mortality**, *see* Deaths
- Mortimer-Filkins test**, 340
- Mothers Against Drunk Driving (MADD)**, 721
- Motivational Enhancement Therapy (MET)**
 less effective in limiting within-treatment drinking, 637
 outcomes of, 124**, 504-12, 631-37
- Motor restraint task**, 562
- mRNA (messenger)**, 16
- Multidimensional Personality Questionnaire (MPQ)**, 200, 222-23, 225
- Multilevel covariance structure analysis (MCA)**, 401-2
- Multiple group structural equation modeling**, 35, 39
- Musculoskeletal complications, and damage from alcohol abuse**, 18
- Myoglobinuria**, 18
- Myopathy in the alcoholic**, 18
- Naloxone, given concomitantly with ethanol**, 258-59
- Naltrexone, to reduce craving following detoxification**, 19
- National Adult Reading Test**, 184
- National Center for Health Statistics (NCHS)**, 456
- National Council on Alcoholism (NCA)**, 469
- 1988 National Health Interview Survey (NHIS)**, 98-99, 105, 191-95
- National Highway Traffic Safety Administration (NHTSA)**, 462, 549
- Fatal Accident Reporting System (FARS)**, 465
- General Estimates System (GES)**, 463
- National Household Survey on Drug Abuse**, 401
- National Institute of Mental Health (NIMH)**, 229
- National Institute on Alcohol Abuse and Alcoholism (NIAAA)**, 20, 89, 98, 289, 471, 631 (*see also* Project MATCH)
 guidelines from, 89, 504
 origin of, 629†
 Research Demonstration Projects from, 524
- National Institute on Drug Abuse (NIDA)**, 120*, 237*, 289
- Collaborative Cocaine Treatment Study (NCCTS)**, 57
- National Household Survey on Drug Abuse**, 401
- 1992 National Longitudinal Alcohol Epidemiologic Survey (NLAES)**, 98-99, 105, 191-95, 664
- National Longitudinal Survey of Youth (NLSY)**, 182, 647-58, 669-80
- National Structured Evaluation (NSE)**, 119*
- Naval Health Research Center (NHRC)**, 272
- Navy Alcohol Treatment Program, reducing lengths of stay in**, 270-79
- Neocortical changes**, 267
- Netherlands Institute for Alcohol and Drugs (NIAD)**, 147
- Neuritis, peripheral**, 29
- Neuronal responsiveness**, 350*
- New York State Cohort (NYSC)**, 293
- Nicotine effects**, *see* Smoking
- N-methyl-D-aspartate (NMDA) receptors, implicated in alcoholic neurologic complications**, 19
- Non-Korsakoff alcoholics**, 180
- Nonspatial conditioned associative-learning task (NSCALT)**, 604, 606
- Normative Aging Study (NAS)**, 712-22
- Nutrition (*see also* Alcohol; Malnutrition)**
 influence on alcohol intake, 26-31
 nutrient degradation, 10
 thiamin deficiencies, 28-29
 unanswered questions about, 30
 and vitamin supplementation, 13
- Nutrition therapy**, 30
 depressing voluntary drinking with, 27-28
- Object Assembly Test**, 563
- "Object" pathway of vision, *versus* "spatial,"** 172
- Occupation and drinking**
 and changes in structure of alcoholic beverage preferences, 112-13
 professionals changing habits more than manual workers and housewives, 112-13
- Ocular-motor control, alcohol-induced reduction in**, 172
- Older drinkers**
 alcohol-related hospitalizations of, 9
 blood alcohol concentrations higher in, 15
 higher rates of cardiomyopathy in, 89, 94
 use of psychotropic drugs, and alcohol interactions, 15
- Onset of drinking**, *see* Drinking
- 1990 Ontario Health Survey (OHS)**, 78-79, 82
 problems with data from, 85, 87
- Ontario Student Drug Use Survey (OSDUS)**, 544-46
- Oppositional defiant disorder (ODD)**, 563
- Organ damage from alcohol abuse**, 16-19
 alcoholic liver disease, 16-17
 cardiovascular effects, 18-19
 endocrine pathology, 17
 gastrointestinal and pancreatic complications, 17
 hematologic disorders, 17-18
 musculoskeletal, renal and dermatologic complications, 18
 neurologic complications, 19
- Osteoporosis and osteonecrosis, alcoholic proneness to**, 15
- Outcome expectancies**
 effect on ad lib alcohol consumption, 198-206
 and theory of planned behavior, 409-19
- Outcomes**, *see* Treatment
- Outcomes Expectancies Scale**, 46, 374
- Out of the labor force (OLF)**, 672
- Outpatient Clinic File (OPC)**, 514-15
- Oversampling**, 99, 524
- Oxandrolone therapy**, 17
- Oxidation (*see also* Alcohol oxidation; Lipid oxidation; Peroxidation)**
 ethanol, to acetaldehyde, via alcohol dehydrogenase (ADH) isoenzymes, 133

- glucose
 impaired, 15, 26
 partial, 27
 with phosphorylation in mitochondria, uncoupled by acetaldehyde, 13
- Paced Auditory Serial Addition Test (PASAT), 181
- Pain relief for alcoholics, and acetaminophen hepatotoxicity, 11
- Pain threshold measures, 604
- Pancreatic complications
 and organ damage from alcohol abuse, 17
 pancreatic insufficiency, 10
- Pancreatitis, *see* Alcoholic pancreatitis
- Parallel analysis, 312
- Parental influences on adolescent drinkers, 32-33
 perception of parental substance abuse, 34
- Paternal absence, 387-98
- Paternal drinking, and development of problem behaviors in boys, 387-98
- Peak weekly alcohol consumption (PAC) level, defining, 135
- Peptic ulceration, 17
- Perceptions (*see also* Media)
 of Japanese junior high school students, 384
 of parental substance abuse, 34
 of treatment institutions by patients, 47
- Perceptual Maze Test, 259-68
- Peripheral neuritis, 29
- Perivenular fibrosis, 16
- Peroxidation
 lipid, 12
 promoted by acetaldehyde, 14
 phospholipid, 16
- Personal Experience Inventory (PEI), 427-28
- Personal History Form (PHF), 525
- Personal interviews, for self-reported alcohol use, 280-85
- Personality Diagnostic Questionnaire-Revised (PDQ-R), 228, 497
- Personality Disorder Examination (PDE), 155, 497
 revised, 495-502
- Personality disorders (PDs), *see* Psychological disorders of alcoholism
- Pharmaceuticals
 blood clearance rate enhanced, 10
 metabolism of increased, 10
- Pharmacologic action of ethanol, 10
 psychoactive properties of, 10
- Phosphatidylethanolamine methyltransferase, 16
- Phospholipids
 peroxidation of, 16
 replenishing *in vitro*, 11
- Physical abuse, *see* Child physical abuse (CPA)
- Picture Arrangement Test, 562
- Pleasurable effects, *see* Alcohol effects
- Point mutations, 133
- Policy issues
 legalization of beer in Iceland, 107-14
 prohibition movement, in Iceland, 107-8
 U.S. Navy treatment programs, 270-79
- Polyenyolphosphatidylcholine, 16
- Polymorphisms, at *ADH2*, *ADH3* and *ALDH2* loci, 135-38
- Porteus Maze Test, 562
- Positive work environment, 44
- Positron emission tomography (PET), for detecting functional cerebral changes, 259-60
- Posttraumatic stress disorder (PTSD), screening for, using Addiction Severity Index (ASI), 56-61
- Predictors
 of alcoholism risk, 485-94
 of alcohol use, 78-88, 314
 of relapse in long-term abstinent (LTA) alcoholics, 640-46
- Prednisolone, improving survival in spontaneous hepatic encephalopathy, 17
- Pregnancy
 abstinence recommended during, 19
 maternal drinking during and after, 292-304, 533-43
- Prenatal alcohol exposure, 292-304
 and family history of alcoholism, 533-43
- Preschool Behaviour Questionnaire, 390
- Prevention, 405
 individualizing, 153
 setting up programs for, 119-20*
- Problem behaviors in boys, link with paternal drinking and paternal absence, 387-98
- Problem-Behavior Theory, 547
- Problem indicators in coworkers, 608-19
- Problems from Substance Use Scale, 515
- Problem solving skills, to improve facility morale, 50
- Processes of Change Inventory, 46
- Prohibitionist approach (*see also* Policy issues)
 poor map for American policy, 238*
- Project MATCH, 5[†], 20, 520
 findings about treatment outcome differences, 124-25**, 503-12, 631-39
- Propylthiouracil, for treating alcoholic hepatitis, 17
- Prosocial Behaviour Questionnaire, 390
- Protective effects
 of aldehyde dehydrogenase (ALDH) isoenzymes, 137-38
 education, 253-54
 of high density lipoproteins (HDL), against atherosclerosis, 18
 of moderate alcohol use, against cardiovascular effects, 18
- Protein, increasing, depresses voluntary drinking, 27-28
- Prothrombin time, elevated, 17
- Proximal outcomes, defined, 371
- Psychiatric comorbidity, 157
- Psychiatric staff, in institutions for treatment, working environment of, 43
- Psychoactive drugs
 in alcoholics with ASPD, 225
 folklore concerning, 619*
 and harm reduction measures, 619-20*
 increased tolerance to, 10
 use correlated with alcohol use, 681-89
- Psychoactive substance use disorder (PSUD), and Executive Cognitive Functioning (ECF), in female adolescents, 560-67
- Psychological disorders of alcoholism, 237-38*, 345
 antisocial personality disorder (ASPD/APD), 39, 233-34, 395, 560
 assessing, 155
 linked with alcoholic depression, 222-26
 attention deficit hyperactivity disorder, 560
 coexisting, 6[†]
 conduct disorder, 560
 depression, and effectiveness of cognitive-behavioral therapy for smokers, 327-35
 depressive-spectrum disease
 alcoholism symptomatic of, 105
 matrilineal heritability of, 106
 inattention/overactivity difficulties, 560
 major depression, 97-106
 in women, 588
 overdiagnosing, 477-78**
 personality disorders (PDs), 499-501

- among treated alcoholics, 227-36
- psychopathy, 560
- secondary depression, 98, 100
- Psychosocial Functioning Inventory, 633
- Psychosocial model of alcoholism, 43-44
- Psychotropic drugs, use by older alcoholics, and alcohol interactions, 15
- PTSD Checklist (PCL), 58
- Pubic hair loss, hallmark of cirrhosis, 17
- Public health aspects of alcohol abuse, 17
- Purine generation, 10
- Quality of Life Enjoyment and Satisfaction Questionnaire (Q-LES-Q), 272-74, 277
- Quantity-frequency (Q-F) measures, 34, 108, 570, 660
- Quantity-Frequency-Variability (QFV), 282
- Random input/output placement, 6†
- Rape, committed under influence of alcohol, 71-72
- Rapid Alcohol Problems Screen (RAPS), 421-25
- Rapid Information Processing (RIP) task, 174-75
- Receiver operating characteristic (ROC) analysis, 555, 557-58
- Reciprocal determinism interaction, 214
- Rectal cancer associated with heavy alcohol consumption, 13
- Redox homeostasis, maintaining, 10
- Relapse prevention (RP) sessions, for couples, 357-70, 591-98
- Relapses
 - administering Disulfiram (Antabuse®) to prevent, 19
 - carrying candies to prevent, 28
 - predicting in long-term abstinent (LTA) alcoholics, 640-46
 - preventing, sessions for couples after male behavioral therapy, 357-70
- Religion, as predictor of current drinking, 665
- Renal complications, and organ damage from alcohol abuse, 18
- Renard Diagnostic Interview, 496
- Replication costs, 508
- Reporting
 - diary, 198
 - electronic interviews, 447-54
 - self, 69, 75, 198, 330, 555-59, 681-89
- Research Council on Problems of Alcohol (RCPA), 629†
- Research Demonstration Projects, 524
- Research Diagnostic Criteria (RDC), 223, 228, 340
- Residential facilities, staff working conditions in, 44
- Respiratory tract cancer associated with heavy alcohol consumption, 13
- Restriction fragment length polymorphism (RFLP), 16
- Retenoids, toxicity of, 13
- Reward system, *see* Alcohol effects
- Rey Auditory Verbal Learning Test, 184-85
- Risk Assessment Battery, 154
- Risk Prediction Scales, 440, 443
- Risk taking, *see* Sexual risk taking
- Robert Wood Johnson Foundation, 570
- Rolling admissions, 154
- Rolling enrollment, 645
- Rural dwelling
 - traditional African rules and regulations, for alcohol consumption, 146-51
 - versus* urban, and changes in structure of alcoholic beverage preferences, 112
- Rutgers Alcohol Problem Index (RAPI), 34, 36-39, 442, 445, 535-41
- S-adenosylmethionine, for treating alcohol-induced liver injury, 17
- Scan order, 259
- Schedule for Affective Disorders and Schizophrenia—Lifetime (SADS-L), 343, 488
- Screening for problem drinking
 - in emergency room populations, performance of different instruments, 420-26
 - in general hospitals, direct or disguised assessment, 698-703
- Seattle Longitudinal Study on Alcohol and Pregnancy, 534
- Secondary depression, *see* Psychological disorders of alcoholism
- Secondary Disabilities study, 620*
- Secondary effects, *see* Alcohol effects
- S.E.D. Laboratories, 556
- Selection bias, *see* Bias
- Self Conscious Scale, 200
- Self-Esteem Inventory (SEI), 537
- Self-help groups, 44
 - Alcoholics Anonymous (AA), 45
 - predictors of participation in, 152-62
- Self-Rating of the Effects of Alcohol (SRE) form, 487
- Self-reporting, 69, 75, 198, 330, 555-59, 681-89
 - mode effects, 280-85
- Semi-Structured Assessment for the Genetics of Alcoholism (SSAGA), 200, 582-83
- Sensation seeking, and alcohol-related injuries, 216-21
- Sensation Seeking Scale, 200, 202, 205, 216, 489
- Serial Seven Test, 259-68
- Serum amylase levels, elevated, 17
- Sexual abuse, *see* Child sexual abuse (CSA)
- Sexual risk taking
 - and alcohol and drug use, 681-89
 - and expectancies of adolescent drinkers, 71-77
- Shibley Institute of Living Vocabulary and Abstraction subscales (SILS), 181-82, 185
- Shocks, administering, 600
- Short-form Alcohol Dependence Data Questionnaire (SADD), 706-9
- Short Michigan Alcoholism Screening Test (SMAST), 387, 389, 496
- Signal detection theory (SDT), 555
- SILS, *see* Shibley Institute of Living Vocabulary and Abstraction subscales (SILS)
- Single photonemission computerized tomography (SPECT), 268
- Situational Confidence Questionnaire (SCQ), 46, 374, 593
- Situational factors, in underage drinking, 207-15, 381-86
 - at bush parties, 544-48
- Skiing injuries, *see* Injuries
- Smoking
 - with alcoholism and depression, cognitive-behavioral therapy for, 327-35
 - effects of prenatal maternal, 292-304
 - lowering activity in anterior cingulate cortex (ACC), 267
 - synergy with alcohol consumption, 255-56
 - and cancer development, 13, 39
 - rationale for, 300
- Sober social drinkers, cognitive functioning in, 180-90
- Sobriety checkpoint programs, 462-68
 - benefits, 464-65
 - costs, 462-64
- Social control theory, 324
- Social drinking, *see* Drinking
- Socialization factors
 - in adolescent drinking
 - parents' norms, 32-33
 - peers' norms, 33, 63-70
 - drinking cultures, 114
 - dynamics of, 39-40
- Social learning theory (SLT), 214

- Social profile, and changes in structure of alcoholic beverage preferences, 110
- Social Provision Scale (SPS), 489
- Social resources, patients with severe lack of, 5-7[†]
- Social settings, underage drinking in, 207-15
- Social Support Questionnaire (SSQ), 489
- Socioeconomic status (SES)
inner-city children, 318
low
controlling for, 387-98
prevalence of psychopathology in, 394
- Solvents, 2E1 activation activity towards, 11
- Sons of male alcoholics (SOMAs), 387-89, 391-95
- Sororities, *see* Student drinkers
- Spatial conditioned associative-learning task (CALT), 600-604
- Spatial contrast sensitivity, effects of moderate BACs on, 163-73
- "Spatial" pathway of vision, *versus* "object," 172
- Spatial Relations test, 182
- Spectrophotometry, 116
- Spielberger Trait Anxiety Scale (STAI), 200
- Spirits, *see* Distilled spirits
- Spontaneous hepatic encephalopathy, prednisolone improving survival in, 17
- Staff working environment, in institutions for treatment, 43-51
high cohesion among staff, 44
psychiatric staff, 43
staff backgrounds and beliefs, 47
- Stages of Change Readiness and Treatment Eagerness Scale, 516
- Standard drink units (SDUs), 288
- Standardized Field Sobriety Tests (SFSTs), 349-50*
- Stress
greater among married drinkers, 110
and workplace drinking, influence of teamwork, 608-18
- Stress-emotional-causal hypothesis, 189
- Stroop Color-Word Interference Test, 184, 562
- Structural equation methodology (SEM), 490
multilevel, 400
- Structural equation modeling, 35, 39, 41
- Structured Clinical Interview for DSM-III-R (SCID)
obtaining diagnoses with, 154-55
SCID-II, 234
- Student drinkers (*see also* Adolescent drinkers; Education)
designated driving among, 549-54
in junior high school in Japan, 381-86
screening, psychometric assessment of SASSI-2, 439-46
socialization of, within Greek societies, 63-70
- Studies (*see also* Animal studies)
by language, Kiswahili, 147
longitudinal, 399-408, 712
follow-up, 89-96, 140-45, 318-26
needed, 237*
into random assignment to treatment approaches, 6[†]
predictors of alcoholism risk, 485-94
replicating, 508
variables, effects of different approaches to consumption measurement, 191-97
- Subjective High Assessment Scale (SHAS), 486
- Substance Abuse Module (SAM), 223
- Substance abuse prevention, *see* Prevention
- Substance Abuse Subtle Screening Inventory (SASSI-2), 439-46
- Substance abuse treatment, *see* Treatment
- Substance use disorder (SUD), 56
- Substance Use Involvement Index, 430
- Suicide
correlation with alcoholism, 144, 455-61
correlation with mental disorder including alcoholism, 144
- Supernutrients, effectiveness in nonhuman primates, 17
- Supervising adolescent drinkers, importance of, 209, 313
- Supervisor's Assessment of Treatment Effectiveness, 272-74
- Suppressor variables, 184
- Survival analysis, 35
- Symbol-Digit Modalities Test, 184
- Symptom Checklist (SCL-90-R), 272
- Symptom Distress Check List (SDCL), 185
- Teamwork, and influence of workplace drinking and stress, 608-18
- Teen Addiction Severity Index (T-ASI), 428
- Telephone surveys, 280-81, 336
- Television, *see* Media
- Temperance values, 107, 112
Mothers Against Drunk Driving (MADD), 721
- Temporal contrast sensitivity, effects of moderate BACs on, 163-73
- Testosterone
plasma, alcohol suppression of, 17
suppressing hepatic ADH activity, 14-15
- Theory of Planned Behavior, 409-19
- Theory of Reasoned Action, 409-10, 415-16
- Therapeutic community work, 120*
- Therapy, *see* Nutrition therapy; Treatment
- Thiamin deficiencies, 28-29
supplementing, 30
- Thoughts About Abstinence Scale, 46, 155
- Thrombocytopenia, 18
- TICS-m test, 186
- Timeline Follow-Back (TLFB) procedure, 155-56, 359, 447-54, 592, 681-89
- Time sequential analyses, 716
- Time series analysis, of alcohol consumption and suicide, 455-61
- Tomography, *see* Positron emission tomography (PET); Single photonemission computerized tomography (SPECT)
- Total Behavior Problem scale, 296
- Total body water (TBW), method for predicting BAC, 350*
- Toxicity mechanisms of alcohol, 10-16
malnutrition, 10
metabolic disorders
from acetaldehyde, 13-14
from alcohol oxidation by alcohol dehydrogenase, 10
from ethanol, carcinogens and vitamin A, 13
gender effects, 14-16
interactions with age, hormones and heredity, 14-16
from microsomal ethanol oxidation, 10-13
role of gastric ADH, 14-16
- Traditional chemical dependency treatment (TCDT) approach, 372
- Tranquilizers, increased concentrations when combined with ethanol, 12-13
- Transferrin levels, *see* Carbohydrate-deficient transferrin (CDT) levels
- Transforming growth factor β , stimulating fibrogenesis, 16
- Trauma, screening for, using Addiction Severity Index (ASI), 56
- Trauma History Questionnaire (THQ), 58-60
- Trauma Scale, 698-702
- Treatment (*see also* Aftercare sessions; Prevention)
disease concept of (DCT), origins and evolution of, 469-76
individualizing, 631-39
innovative approaches, effectiveness of, 120-21*
inpatient
move to abolish, 5-7[†]
move to shorten stay in U.S. Navy program, 270-79
integrated programs, 120*
length, 270-79
motives for seeking, 517
multimodal, 120*
outcomes

- better for patients in inpatient treatment, 5[†]
 for couples in relapse prevention sessions, 357-70
 differences between three types of, 503-12, 631-39
 Cognitive Behavioral Coping Skills Therapy (CBT),
 124**, 504-12, 631-37
 Motivational Enhancement Therapy (MET), 124**,
 504-12, 631-37
 Twelve-Step Facilitation Therapy (TSF), 124**, 504-12,
 631-37
 for Hispanics *versus* non-Hispanic whites, 286-91
 influence of outpatient treatment and 12-step group involve-
 ment on, 513-22
 proximal, variables during 12-step and cognitive-behavioral
 treatment, 371-80
- outpatient
 detoxification, successes in, 6[†]
 intensive, participation in aftercare sessions and self-help
 groups following, 152
 and 12-step group involvement, 513-22
 residential, in therapeutic community work, 120*
 resistance to, patients with high levels of, 6[†]
 substance abuse, process models for evaluating, 371-80
 utilization rates, Hispanics *versus* non-Hispanic
 whites, 286-91
- Treatment Experiences Form, 46
 Treatment institutions, 43-51
 characteristics of, 43
 patients' discharge outcomes, 48
 patients' improvement during treatment, 44
 patients' involvement in, 44, 47-48
 patients' perceptions of, 47
 staff working environment in, 43-51
- Treatment matching, 631-39
 Treatment Outcome Prospective Study (TOPS), 515
 Treatment Services Interview (TSI), 525
 Treatment Services Review (TSR), 155-56
- Trends
 in alcohol consumption patterns, among whites, blacks and His-
 panics, 659-68
 changes in structure of alcoholic beverage preferences, 109-10
 decreases in consumption of distilled spirits, 107
 decreases in overall alcohol consumption, 19-20, 107
 move to abolish inpatient treatment, 5-7[†]
 relative increases in consumption of lower alcohol level bever-
 ages, 107
- Tridimensional Personality Questionnaire, 489
 Tumor necrosis factor α , stimulating fibrogenesis, 16
 TWEAK, 420-24, 698-99, 702
 Twelve-Step Facilitation Therapy (TSF), outcomes of, 124**,
 504-12, 631-37
- 12-step programs, 371-80
 and outpatient treatment, 513-22
 working the steps, 696
- 2E1 activation activity, 11
 Type I censoring, 35
 Type I drinkers, adolescent, 32-33
 Type I errors, 65, 431, 563
 Type II drinkers, 395
 adolescent, 32-33
 Type II errors, 563
- Underage drinking, social settings and situations, 207-15,
 381-86
- Underemployment, and alcohol misuse, 669-80
 Underreporting drinking, 109, 150, 346
 Understanding of Alcoholism Scale, 373
 Understanding of Substance Abuse Scale, Short Form of, 45
- 1971 Uniform Alcoholism and Intoxication Treatment Act, 472
 United Nations, Food and Agriculture Organization (FAO), 150
 U.S. Preventive Services Task Force, guidelines from, 89
 U.S. standard drinks (USSD), 181-83, 185
 Urban *versus* rural dwelling, and changes in structure of alcoholic
 beverage preferences, 112
- Vaillant's Core City sample, 143
 Vascular spiders, hallmark of cirrhosis, 17
 Vegetarian diet, and alcohol abuse, 29
 Verbal intelligence, 182
 Vestibulo-ocular reflex (VOR), 724-26, 728-29
 Veterans Affairs Beneficiary Identification and Record Locator
 Subsystem, 90
 Veterans Affairs hospitals, *see* Department of Veterans Affairs
 (VA) Medical Centers
 Veterans Affairs nationwide outpatient clinic (OPC) file, 46
 Veterans Affairs Patient Treatment File (PTF), 373, 514
 Veterans, death rates from alcohol-related liver diseases, 9
 Viral hepatitis, 16
 Visual acuity, 163
 Vitamin deficiencies, 28
 vitamin A, and cancer associated with heavy alcohol consump-
 tion, 13
 vitamin B, 10
- Vulnerability
 social, 614-15
 of women, 14
- Wanzuki, 149
 Wasukuma tribe, 146-51
 Wave analysis, 312-16
 Ways of Coping Scale (WOC), 489-90
 Wechsler Adult Intelligence Scale (WAIS), Vocabulary, Similari-
 ties and Comprehension, 184
 Wechsler Adult Intelligence Scale—Revised (WAIS-R), 182-83
 short form of, 600
 Wechsler Intelligence Scale for Children—Revised (WISC-R), 562
 Wechsler Memory Scale, 183-84
 Logical Memory, 183
 Mental Control, 183
 Visual Reproduction (Immediate and Delayed), 183
- Wernicke-Korsakoff's syndrome, 29
 Wisconsin Card Sorting Test (WCST), short form, 181
- Women
 carbohydrate-deficient transferrin (CDT) levels in, 555-59
 clinical course of alcohol-related problems in, 581-90
 disapproval of female drunkenness, 40
 genetic factors stronger in, comorbidity indicating common un-
 derlying cause, 97-98
 increased vulnerability to alcohol, 14
 making women aware of, 14
 maternal drinking, during and after pregnancy, 292-304, 533-43
 matrilineal heritability, of depressive-spectrum disease, 106
 menstrual cycle disorders in, 14-15
 hallmark of cirrhosis, 17
 during pregnancy, recommending abstinence, 19
 social drinking among, 14
 trend towards complete abstention, 660-62
 young
 psychoactive substance use disorder (PSUD), and Executive
 Cognitive Functioning (ECF), 560-67
 socialization regarding alcohol use, 33
- Woodlawn study, 318-26
 Woodruff principle, 477-78**
 Work Environment Scale (WES), 44
 subscales to, 46

- Workplace morale
and drinking, influence of teamwork, 608-18
in institutions for treatment, 43-51
- Work Stress Questionnaire, 489
- World Drink Trends, 1993*, 246
- World Health Organization (WHO), 185
alcohol questionnaire, 343
conference on population levels of alcohol consumption, 250
- definitions, 700
modification of PDE, 228
World Health Statistics Annuals, 246-47
- Xenobiotics, activated to toxic metabolites, 11
- Yale Plan Clinics, 472-73